Genomeditech (Shanghai) Co., Ltd.
Order: +86 021-50432825/50432826
Email: service@genomeditech.com

产品手册

H_IL31RA CHO-K1 Cell Line

货号: GM-C25992

版本号：4.0.26.1015

规格参数

产品描述	H_IL31RA CHO-K1 Cell Line 采用慢病毒技术构建，为单克隆稳定细胞株，组成型表达人 IL31RA。
数量	5E6 Cells per vial,1 mL
产品形式	1 支冻存细胞
储运条件	干冰运输， －196° C 以下长期储存
靶点	Human_IL31RA
转录本	Q8NI17-1(AA Met 1 - Cys 553)
宿主细胞	CHO-K1
复苏培养基	F12K+10% FBS+1% P.S
生长培养基	F12K+10% FBS+1% P.S+4 μg/mL Puromycin
培养备注	无
细胞冻存液	90% FBS+10% DMSO
生长特性	贴壁
培养条件	37°C, 5% CO_2
支原体检测	该细胞系已通过支原体检测，结果为阴性。
生物安全等级	Biosafety Level 2
备注	建议尽早扩增该细胞，并冻存至少 10 支，以便后续使用。


本公司产品仅供科研用途，严禁用于人体治疗！

实验材料与仪器

名称	Manufacturer/Catalogue No.
F12K	BOSTER/PYG0036
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/GM-040401
Anti-IL31RA hIgG2 Antibody(Nemolizumab)	Genomeditech/GM-50871AB
Countess 3 Automated Cell Counter	ThermoFisher/A49891
NovoCyte Flow Cytometer	Agilent/NovoCyte 2060R

验证结果

表达



	SampleID	Geometric Mean : R675-H
■	CHO-K1 anti-IL31RA+APC-2nd	611
■	CHO-K1 H_IL31RA H_IgG+APC-2nd	735
■	CHO-K1 H_IL31RA anti-IL31RA+APC-2nd	380804

IL31RA 流式表达验证。采用流式细胞术，用 Anti-IL31RA hIgG2 抗体（GM-50871AB）检测 H_IL31RA CHO-K1 细胞系（GM-C25992）的 IL31RA 表达。

细胞复苏

复苏培养基：F12K+10% FBS+1% P.S

为确保最高存活率，应在收到冻存细胞后立即解冻并复苏培养。如果在收到细胞后需要继续储存，将其置于液氮罐中，严禁储存在-70° C，因为在-70° C 下储存会导致活性丧失。

a) 37℃水浴锅预热复苏培养基，加入预热后的复苏培养基 5 mL 至 15 mL 离心管。

b) 从液氮中取出冻存细胞并迅速放入 37℃恒温水浴锅，将细胞液面浸至水面以下轻轻摇动解冻，直到刚刚融化（通常 2-3 分钟）。

c) 用 70%乙醇擦拭冻存管外部以降低污染的几率。在生物安全柜或超净台中将冻存管中的细胞悬液转移到步骤 a）的离心管中，轻轻混匀，176 × g，离心 3 min，使细胞沉淀，弃上清。

d) 使用 1 mL 复苏培养基重悬，可取出部分使用台盼蓝染色计数活细胞，活细胞≥ 3E6 cells/mL。

e) 通过补加复苏培养基的形式，调整活细胞密度到 2-3E5 cells/mL，根据细胞悬液总体积，将细胞接种到合适的培养皿中。

细胞冻存

细胞冻存液：90% FBS+10% DMSO

a) 使用 176 × g，3 min 离心收集细胞。

b) 使用预冷细胞冻存液（90% FBS + 10% DMSO）重悬细胞，细胞密度调整为 5E6 cells/mL，每管 1 mL 分装到细胞冻存管中。拧紧盖子，适当标记后，将冻存管置于梯度降温盒中，-80℃下保存至少 1 天，尽快转移至液氮中。

细胞传代

生长培养基： F12K+10% FBS+1% P.S+4 μg/mL Puromycin

细胞复苏后的 1 至 2 代，使用复苏培养基，待细胞状态稳定后，再更换为含有抗生素的生长培养基。

a) 此细胞呈梭状，贴壁生长。培养箱中孵育 16-24 h 后，镜下观察细胞贴壁情况。当细胞密度大于 60%，即可进行细胞传代。

b) 将皿或培养瓶中的培养液弃去，10 cm 皿加 2 mL PBS 润洗 1 次。

c) 弃 PBS，加 1 mL 0.25% Trypsin-EDTA 消化液，37° C 消化 2-3 min，显微镜下观察。

d) 待细胞变圆，细胞间隙明显，部分细胞刚开始脱离瓶壁时，加 2 mL 左右生长培养基混匀终止消化，将细胞小心吹打下来，176 × g 室温离心 3 min。

e) 弃上清，细胞沉淀用生长培养基重悬，根据传代前细胞密度分盘（根据培养皿面积和细胞密度计算，传代后细胞密度为 20-30%）。

细胞传代比例为 1:4-1:5，2-3 天传代。

培养备注

a) 细胞状态稳定后，传代后死细胞会变少，细胞生长速度趋于稳定，细胞形态均匀，胞体健壮。

序列

IL31RA Q8NI17-1(ΔICD)

MMWTWALWMLPSLCKFSLAALPAKPENISCVYYYRKNLTCTWSPGKETSYTQYTVKRTYAFGEKHDNCTTNS
STSENRASCSFFLPRITIPDNYTIEVEAENGDGVIKSHMTYWRLENIAKTEPPKIFRVKPVLGIKRMIQIEWIKPELA
PVSSDLKYTLRFRTVNSTSWMEVNFAKNRKDKNQTYNLTGLQPFTEYVIALRCAVKESKFWSDWSQEKMGMTE
EEAPCGLELWRVLKPAEADGRRPVRLLWKKARGAPVLEKTLGYNIWYYPESNTNLTETMNTTNQQLELHLGGE
SFWVSMISYNSLGKSPVATLRIPAIQEKSFQCIEVMQACVAEDQLVVKWQSSALDVNTWMIEWFPDVDSEPTTLS
WESVSQATNWTIQQDKLKPFWCYNISVYPMLHDKVGEPYSIQAYAKEGVPSEGPETKVENIGVKTVTITWKEIPK
SERKGIICNYTIFYQAEGGKGFSKTVNSSILQYGLESLKRKTSYIVQVMASTSAGGTNGTSINFKTLSFSVFEIILITS
LIGGGLLILIILTVAYGLKKPNKLTHLC

相关产品

OX40:OX40L	
H_OX40 Reporter Cell Line	H_OX40 Reporter DDX35TM Cell Line
Cynomolgus_OX40 HEK-293 Cell Line	Cynomolgus_OX40L CHO-K1 Cell Line
H_OX40 CHO-K1 Cell Line	H_OX40L CHO-K1 Cell Line
H_OX40L HEK-293 Cell Line	Mouse_OX40L CHO-K1 Cell Line
Rat_OX40L CHO-K1 Cell Line	
Anti-H_OX40 hIgG2 Antibody(Ivuxolimab)	Anti-OX40L hIgG1 Reference Antibody(Oxebio)
Anti-OX40L hIgG4 Antibody(Amlitelimab)	Anti-OX40L hIgG4 Reference Antibody(Amlbio)
Biotinylated Human OX40L Protein; His-Avi Tag	Cynomolgus OX40 Protein; His Tag
Cynomolgus OX40L Protein; His Tag	Cynomolgus OX40L Protein; mFc Tag
Human OX40 Protein; hFc Tag	Human OX40 Protein; His Tag
Human OX40L Protein; His Tag	Human OX40L Protein; mFc Tag
IL-4/IL-13	
IL-4 Reporter Cell Line	IL-4/IL-13 Reporter 293 Cell Line
IL-4/IL-13 Reporter 293 DDX35TM Cell Line	Cynomolgus_IL4R CHO-K1 Cell Line
H_IL4R CHO-K1 Cell Line	H_IL4R CHO-K1 Cell Line (Low Expression)
H_IL4R HEK-293 Cell Line	Mouse_IL4R CHO-K1 Cell Line
Anti-IL13 hIgG4 Reference Antibody (Lebribio)	Anti-IL-4R hIgG1 Antibody(12B5)
Anti-IL4R hIgG4 Antibody(Dupilumab)	Anti-IL4R hIgG4 Reference Antibody (Dupbio)
Anti-IL-4R×IL31 hIgG4 Reference Antibody (PRO2198)	Anti-Mouse IL-4RA mIgG1 Antibody
Biotinylated Human IL-4R alpha Protein; Avi-His Tag	Cynomolgus IL-13 Protein; His Tag
Cynomolgus IL-13RA1 Protein; hFc Tag	Cynomolgus IL-13RA1 Protein; His Tag
Cynomolgus IL-4 Protein; His Tag	Cynomolgus IL-4R alpha Protein; His Tag
Human IL-13 Protein; His Tag	Human IL-13RA1 Protein; hFc Tag
Human IL-13RA1 Protein; His Tag	Human IL-4 Protein; His Tag
Human IL-4R alpha Protein; hFc Tag	Human IL-4R alpha Protein; His Tag
Human IL-4R alpha Protein; mFc Tag	Mouse IL-13 Protein; His Tag

Mouse IL-4R alpha Protein; His Tag	Rat IL-4R alpha Protein; His Tag
IL-31	
Cynomolgus_IL-31RA OSMR Reporter Baf3 Cell Line	H_IL-31 Reporter Cell Line
H_IL-31 Reporter DDX35TM Cell Line	Cynomolgus_IL31RA CHO-K1 Cell Line
H_IL31RA HEK-293 Cell Line	H_IL-31RA OSMR Baf3 Cell Line
H_IL-31RA OSMR CHO-K1 Cell Line	
Anti-IL31 hIgG1 Antibody(mAb33)	Anti-IL-31 hIgG4 Reference Antibody (BMS-981164)
Anti-IL31RA hIgG1 Antibody(NA633)	Anti-IL31RA hIgG2 Antibody(Nemolizumab)
Anti-IL31RA hIgG2 Reference Antibody (Nemobio)	Anti-IL-4R×IL31 hIgG4 Reference Antibody (PRO2198)
Anti-OSMR hIgG4 Antibody(Vixarelimab)	Anti-OSMR hIgG4 Reference Antibody (Vixabio)
Cynomolgus IL-31 Protein; His Tag	Cynomolgus IL-31RA Protein; hFc Tag
Human IL-31 Protein; hFc Tag	Human IL-31 Protein; His Tag
Human IL-31RA Protein; hFc Tag	
MRGPRX2	
H_MRGPRX2 Gq Reporter CHO-K1 Cell Line	H_MRGPRX2 Gqi5 Reporter CHO-K1 Cell Line
Tango-H_MRGPRX2 CHO-K1 Cell Line	Cynomolgus_MRGPRX2 CHO-K1 Cell Line
Cynomolgus_MRGPRX2 HEK-293 Cell Line	Flag-Mouse_Mrgprb2 CHO-K1 Cell Line
Flag-Rat_Mrgprb3 HEK-293 Cell Line	H_MRGPRX2 CHO-K1 Cell Line
H_MRGPRX2 HEK-293 Cell Line	H_MRGPRX2 HMC-1 Cell Line
H_MRGPRX2 RBL-2H3 Cell Line	
STAT6	
H_IL4R STAT6-HiBiT-KI LgBiT Jurkat Cell Line	

使用许可协议

凡购买及使用本细胞系产品，即表明使用者自愿接受并遵守以下相关使用政策：

- 本细胞系产品限于科研用途，不得被利用于任何商业用途。
- 本产品严禁用于人类或动物疾病诊治，也不得直接用于人体相关实验。
- 用户及为其利益服务的第三方承包商仅可在约定科研范围内使用本材料及其子代，不得进行修饰，亦不得向任何其他实体（包括关联机构）分发、销售、转让或以其他方式提供吉满生物材料。
- 如需将本产品用于本声明范围以外的用途，须事先获得吉满生物科技（上海）有限公司的书面许可，详情请联系吉满生物科技（上海）有限公司。