

## Mouse VEGFR2 Protein; His Tag

## **Product Information**

Product Name	Mouse VEGFR2 Protein; His Tag
Storage temp.	Store at $\leq$ -70°C, stable for 6 months after receipt. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Catalog# / Size	GM-87864RP-100 / 100 μg GM-87864RP-1000 / 1 mg

## **Protein Information**

Alternative Names	KDR,CD309,FLK1,VEGF R2
Source	Mouse VEGFR2 Protein; His Tag (GM-87864RP) is expressed from human 293
	cells (HEK-293). It contains AA Ala 20 - Glu 762 (Accession # P35918-1).
	This protein carries a His tag at the C-terminus.
Purity	> 90% as determined by SDS-PAGE
Endotoxin	$< 1 \text{ EU/}\mu g$ , determined by LAL gel clotting assay
Predicted Mol Mass	82.5 KDa
Formulation	Supplied as a 0.2 $\mu$ m filtered solution of PBS, pH7.2-7.4.
Description	VEGFR2 (Vascular Endothelial Growth Factor Receptor 2) is a key tyrosine
	kinase receptor in vascular endothelial cells. It binds to VEGF, activating
	pathways like PI3K/AKT and MAPK to regulate angiogenesis, vascular
	permeability, cell proliferation, and migration. VEGFR2 is crucial for embryonic
	development and tumor growth, with abnormal expression linked to tumor
	angiogenesis and invasion, making it a key anti-angiogenesis target.
	Composed of extracellular, transmembrane, and intracellular domains, VEGFR2
	shows abnormal expression in conditions like cancer and ocular diseases.
	Targeted inhibitors, such as tyrosine kinase inhibitors and antibody drugs, are
	widely used in clinical therapies, marking VEGFR2 as a focus for cancer and anti-
	angiogenesis drug research.
Description	kinase receptor in vascular endothelial cells. It binds to VEGF, activating pathways like PI3K/AKT and MAPK to regulate angiogenesis, vascular permeability, cell proliferation, and migration. VEGFR2 is crucial for embryonic development and tumor growth, with abnormal expression linked to tumor angiogenesis and invasion, making it a key anti-angiogenesis target. Composed of extracellular, transmembrane, and intracellular domains, VEGFR2 shows abnormal expression in conditions like cancer and ocular diseases. Targeted inhibitors, such as tyrosine kinase inhibitors and antibody drugs, are widely used in clinical therapies, marking VEGFR2 as a focus for cancer and anti-




**SDS-PAGE** 



On SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.

## **Bioactivity-ELISA**



Mouse VEGFR2 Protein; His Tag (Catalog # GM-87864RP) was immobilized at 2 µg/ml (100 µL/well). Increasing concentrations of Anti-mouse VEGFR-2 mIgG2a Antibody(DC101) (Catalog # GM-87892MAB) were added.