

Human TNFR2 Protein; His Tag

Product Information


Product Name	Human TNFR2 Protein; His Tag
Storage temp	Store at $\leq -70^{\circ}\text{C}$, stable for 6 months after receipt. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Catalog# / Size	GM-88213RP-100 / 100 μg GM-88213RP-1000 / 1 mg

Protein Information

Alternative Names	TNFRSF1B, CD120b, TBPII, TNF-R-II, TNF-R75, TNFBR, TNFR1B, TNFR2
Source	Human TNFR2 Protein; His Tag (GM-88213RP) is expressed from human 293 cells (HEK-293). It contains AA Leu 23 - Asp 257 (Accession # P20333-1). This protein carries a His tag at the C-terminus.
Purity	> 95% as determined by SDS-PAGE
Endotoxin	< 1 EU/ μg , determined by LAL gel clotting assay
Predicted Mol Mass	26.0 KDa
Formulation	Supplied as a 0.2 μm filtered solution of PBS, pH7.2-7.4.
Description	<p>TNF receptors 1 and 2 (TNFR1 and TNFR2) mediate the effects of tumor necrosis factor (TNF). TNFR1, found on most nucleated cells, has a death domain and can trigger apoptosis or NF-κB/MAPK-mediated survival signals, activated by both soluble and membrane-bound TNF. TNFR2, mainly on immune and endothelial cells (lacks a death domain), promotes cell survival and regulatory T cell function via NF-κB and related pathways, often in response to membrane-bound TNF. The two receptors can form heterotrimers and modulate each other's signaling, reflecting TNF biology's complexity.</p> <p>TNF-α signals primarily through its receptors TNFR1 (p55) and TNFR2 (p75), triggering multiple pathways. TNFR1 activation commonly leads to NF-κB and MAPK signaling, promoting transcription of inflammatory genes, survival factors, and sometimes apoptosis via caspase cascades. TNFR2 mainly modulates immune cell activation and can enhance TNF-α-driven responses. Cross-talk with other cytokines shapes the intensity and duration of inflammation. Therapeutically, TNF-α inhibitors are used to treat autoimmune diseases, but neutralization must balance infection risk and immune suppression.</p>


Version:4.0

SDS-PAGE




On SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA




Human TNFR2 Protein; His Tag (Catalog # GM-88213RP) was immobilized at 2 μ g/ml (100 μ L/well). Increasing concentrations of Human TNF alpha Protein; hFc Tag (Catalog # GM-88210RP) were added.




Human TNFR2 Protein; His Tag (Catalog # GM-88213RP) was immobilized at 2 μ g/ml (100 μ L/well). Increasing concentrations of Anti-H_TNFRSF1B(TNFR2) hIgG1 Antibody(UC2.3.8) (Catalog # GM-49245AB) were added.

Bioactivity-ELISA




Human TNFR2 Protein; His Tag (Catalog # GM-88213RP) was immobilized at 2 µg/ml (100 µL/well). Increasing concentrations of Anti-H_TNFRSF1B(TNFR2) hIgG1 Antibody (Catalog # GM-49248AB) were added.

Bioactivity-ELISA




Human TNFR2 Protein; His Tag (Catalog # GM-88213RP) was immobilized at 2 µg/ml (100 µL/well). Increasing concentrations of Anti-H_TNFR2 hIgG1 Antibody(1H10) (Catalog # GM-59476AB) were added.

Bioactivity-ELISA



Human TNFR2 Protein; His Tag (Catalog # GM-88213RP) was immobilized at 2 µg/ml (100 µL/well). Increasing concentrations of Anti-H_TNFR2 hIgG2 Antibody(1) (Catalog # GM-59477AB) were added.

Bioactivity-ELISA



Human TNFR2 Protein; His Tag (Catalog # GM-88213RP) was immobilized at 2 µg/ml (100 µL/well). Increasing concentrations of Anti-H_TNFR2 hIgG2 Antibody(13) (Catalog # GM-59478AB) were added.